

MIA3 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A17589	Rabbit	1 mg/ml	
Applications	WB,ELISA		
Reactivity	Human,Mouse		
Dilution	WB 1:500-2000 ELISA 1:5000-20000		
Storage	-20°C/1 year		
Specificity	MIA3 Polyclonal Antibody detects endogenous levels of protein.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthesized peptide derived from human protein . at AA range: 1110-1190		
Uniprot No	Q5JRA6		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	<p>domain:Although 2 transmembrane domains are predicted, PubMed:19269366 showed that it only contains one transmembrane domain. The other predicted transmembrane region is probably a hairpin-type region embedded into the membrane, which does not cross the membrane. It is unclear which of the 2 predicted transmembrane regions is the transmembrane or the hairpin-type region.,domain:The proline-rich region (PRD) mediates the interaction with COPII coat subunits Sec23/24.,function:Required for collagen VII (COL7A1) secretion by loading COL7A1 into transport carriers. May participate in cargo loading of COL7A1 at endoplasmic reticulum exit sites by binding to COPII coat subunits Sec23/24 and guiding SH3-bound COL7A1 into a growing carrier. Does not play a role in global protein secretion and is apparently specific to COL7A1 cargo loading. However, it may participate in secretion of other proteins in cells that do not secrete COL7A1.,similarity:Belongs to the MIA/OTOR family. Tango1 subfamily.,similarity:Contains 1 SH3 domain.,subcellular location:Localizes at endoplasmic reticulum exit sites. After loading of COL7A1 into transport carriers, it is not incorporated into COPII carriers and remains in the endoplasmic reticulum membrane.,subunit:Interacts (via SH3 domain) with COL7A1. Associates with the COPII coat subunits Sec23/Sec24.,tissue specificity:Broadly expressed, except in bone marrow and peripheral blood mononuclear cells. Down-regulated in melanoma tissue..</p>		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
