

## NMDAε2 (phospho Ser1303) rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A18450	Rabbit	1 mg/ml	165959
<b>Applications</b>	WB,ELISA		
<b>Reactivity</b>	Human,Mouse,Rat		
<b>Dilution</b>	WB: 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.		
<b>Storage</b>	-20°C/1 year		
<b>Specificity</b>	Phospho-NMDAε2 (S1303) Polyclonal Antibody detects endogenous levels of NMDAε2 protein only when phosphorylated at S1303.		
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human GRIN2B around the phosphorylation site of Ser1303. AA range:1269-1318		
<b>Uniprot No</b>	Q13224		
<b>Alternative names</b>	GRIN2B; NMDAR2B; Glutamate [NMDA] receptor subunit epsilon-2; N-methyl D-aspartate receptor subtype 2B; NMDAR2B; NR2B; N-methyl-D-aspartate receptor subunit 3; NR3; hNR3		
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Conjugation</b>			
<b>Background</b>	<p>glutamate ionotropic receptor NMDA type subunit 2B(GRIN2B) Homo sapiens N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA receptor channel has been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of three different subunits: NR1 (GRIN1), NR2 (GRIN2A, GRIN2B, GRIN2C, or GRIN2D) and NR3 (GRIN3A or GRIN3B). The NR2 subunit acts as the agonist binding site for glutamate. This receptor is the predominant excitatory neurotransmitter receptor in the mammalian brain. [provided by RefSeq, Jul 2008],</p>		
<b>Other</b>	GRIN2B, Glutamate [NMDA] receptor subunit epsilon-2		

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---