

## Pin1 (phospho Ser16) rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A19857	Rabbit	1 mg/ml	18243

<b>Applications</b>	WB,IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat,Monkey
<b>Dilution</b>	WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. ELISA: 1:20000. Not yet tested in other applications.
<b>Storage</b>	-20°C/1 year
<b>Specificity</b>	Phospho-Pin1 (S16) Polyclonal Antibody detects endogenous levels of Pin1 protein only when phosphorylated at S16.
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Pin1 around the phosphorylation site of Ser16. AA range:1-50
<b>Uniprot No</b>	Q13526
<b>Alternative names</b>	PIN1; Peptidyl-prolyl cis-trans isomerase NIMA-interacting 1; Peptidyl-prolyl cis-trans isomerase Pin1; PPIase Pin1; Rotamase Pin1
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Conjugation</b>	
<b>Background</b>	peptidylprolyl cis/trans isomerase, NIMA-interacting 1(PIN1) Homo sapiens Peptidyl-prolyl cis/trans isomerases (PPIases) catalyze the cis/trans isomerization of peptidyl-prolyl peptide bonds. This gene encodes one of the PPIases, which specifically binds to phosphorylated ser/thr-pro motifs to catalytically regulate the post-phosphorylation conformation of its substrates. The conformational regulation catalyzed by this PPIase has a profound impact on key proteins involved in the regulation of cell growth, genotoxic and other stress responses, the immune response, induction and maintenance of pluripotency, germ cell development, neuronal differentiation, and survival. This enzyme also plays a key role in the pathogenesis of Alzheimer's disease and many cancers. Multiple alternatively spliced transcript variants have been found for this gene.[provided by RefSeq, Jun 2011],
<b>Other</b>	PIN1, Peptidyl-prolyl cis-trans isomerase NIMA-interacting 1

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---