

PPIG rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A20130	Rabbit	1 mg/ml	

Applications	WB,ELISA
Reactivity	Human,Rat,Mouse
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage	-20°C/1 year
Specificity	PPIG Polyclonal Antibody detects endogenous levels of protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthesized peptide derived from human protein . at AA range: 290-370
Uniprot No	Q13427
Alternative names	
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	IgG
Conjugation	
Background	<p>catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,domain:The RS domain is required for the interaction with the phosphorylated C-terminal domain of RNA polymerase II.,enzyme regulation:Cyclosporin A (CsA)-sensitive.,function:PPIases accelerate the folding of proteins.,function:PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,function:PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be implicated in the folding, transport, and assembly of proteins. May play an important role in the regulation of pre-mRNA splicing.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the cyclophilin-type PPIase family.,similarity:Contains 1 PPIase cyclophilin-type domain.,subcellular location:Colocalizes with RNA splicing factors at nuclear speckles.,subunit:Interacts with CLK1, PNN and with the phosphorylated C-terminal domain of RNA polymerase II.,tissue specificity:Ubiquitous.,</p>
Other	PPIG, Peptidyl-prolyl cis-trans isomerase G (PPIase G) (Peptidyl-prolyl isomerase G) (EC 5.2.1.8) (CASP10) (Clk-associating RS-cyclophilin) (CARS-Cyp) (CARS-cyclophilin) (SR-cyclophilin) (SR-cyp) (SRcyp) (Cyclophilin G) (Rotamase G)

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
