

## SLAP1 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A21586	Rabbit	1 mg/ml	

<b>Applications</b>	WB,ELISA
<b>Reactivity</b>	Human,Rat,Mouse
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Storage</b>	-20°C/1 year
<b>Specificity</b>	SLAP1 Polyclonal Antibody detects endogenous levels of protein.
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Uniprot No</b>	Q13239
<b>Alternative names</b>	
<b>Form</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Conjugation</b>	
<b>Background</b>	<p>domain:The C-terminal domain is essential for the homodimerization and the interaction with CBL. While the interaction with CBL is apparently mediated via the hydrophobic region of this domain, the highly charged region is apparently required for the homodimerization.,function:Adapter protein, which negatively regulates T-cell receptor (TCR) signaling. Inhibits T-cell antigen-receptor induced activation of nuclear factor of activated T-cells. Involved in the negative regulation of positive selection and mitosis of T-cells. May act by linking signaling proteins such as ZAP70 with CBL, leading to a CBL dependent degradation of signaling proteins.,induction:By all-trans retinoic acid (ATRA). Induction is indirect and is mediated through other proteins.,PTM:Phosphorylated.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 SH3 domain.,subcellular location:Colocalizes with endosomes.,subunit:Interacts with EPHA2, VAV1, LCP2 and PDGFRB (By similarity). Homodimer. Homodimerization and interaction with phosphorylated CBL occurs via its C-terminal domain. Interacts with phosphorylated proteins ZAP70, CD3Z, SYK and LAT via its SH2 domain.,tissue specificity:Expressed in lung and fetal brain. Weakly expressed in heart, adult brain, placenta, liver, skeletal muscle, kidney and pancreas.,</p>

**Other** ~~SLA SLAP SLAP1, Src-like-adaptor (Src-like-adaptor protein 1) (SLAP-1)~~

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---