

TLR13 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A22623	Rabbit	1 mg/ml	

Applications	IHC
Reactivity	Human,Mouse,Rat
Dilution	IHC: 1:200-500
Storage	-20°C/1 year
Specificity	The antibody detects endogenous TLR13 protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Recombinant Protein of TLR13
Uniprot No	
Alternative names	
Form	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Clonality	Polyclonal
Isotype	IgG
Conjugation	
Background	The TLRs form a family of ligand-binding proteins that engage a variety of microbial products, that on binding, activate signaling cascades leading to the synthesis of proinflammatory molecules. TLR13 is found in mouse and participates in the innate immune response to microbial agents. It acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response.
Other	TLR13, Toll-like receptor 13
Product Images:	

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.