

Cleaved-Tumstatin (P1426) rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A12676 | Rabbit | 1 mg/ml | 161813 |
| Applications | WB,ELISA | | |
| Reactivity | Human,Monkey | | |
| Dilution | WB: 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications. | | |
| Storage | -20°C/1 year | | |
| Specificity | Cleaved-Tumstatin (P1426) Polyclonal Antibody detects endogenous levels of fragment of activated Tumstatin protein resulting from cleavage adjacent to P1426. | | |
| Source / Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. | | |
| Immunogen | The antiserum was produced against synthesized peptide derived from human Collagen IV alpha3. AA range:1407-1456 | | |
| Uniprot No | Q01955 | | |
| Alternative names | COL4A3; Collagen alpha-3(IV) chain; Goodpasture antigen | | |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | | |
| Clonality | Polyclonal | | |
| Isotype | IgG | | |
| Conjugation | | | |
| Background | <p>collagen type IV alpha 3 chain(COL4A3) Homo sapiens Type IV collagen, the major structural component of basement membranes, is a multimeric protein composed of 3 alpha subunits. These subunits are encoded by 6 different genes, alpha 1 through alpha 6, each of which can form a triple helix structure with 2 other subunits to form type IV collagen. This gene encodes alpha 3. In the Goodpasture syndrome, autoantibodies bind to the collagen molecules in the basement membranes of alveoli and glomeruli. The epitopes that elicit these autoantibodies are localized largely to the non-collagenous C-terminal domain of the protein. A specific kinase phosphorylates amino acids in this same C-terminal region and the expression of this kinase is upregulated during pathogenesis. This gene is also linked to an autosomal recessive form of Alport syndrome. The mutations contributing to this syndrome are also located within the exons that encode this C-terminal r</p> | | |
| Other | COL4A3, Collagen alpha-3(IV) chain | | |

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
