

## CRLF3 rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|--------------|---------|-----------------|---------------|
| A13067       | Rabbit  | 1 mg/ml         |               |

|                              |  |
|------------------------------|--|
| <b>Applications</b>          | WB,ELISA   |
| <b>Reactivity</b>            | Human,Mouse  |
| <b>Dilution</b>              | WB 1:500-2000 ELISA 1:5000-20000   |
| <b>Storage</b>               | -20°C/1 year   |
| <b>Specificity</b>           | CRLF3 Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen</b>             | Synthesized peptide derived from human protein . at AA range: 270-350  |
| <b>Uniprot No</b>            | Q8IUI8   |
| <b>Alternative names</b>     |  |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Clonality</b>             | Polyclonal   |
| <b>Isotype</b>               | IgG  |
| <b>Conjugation</b>           |  |
| <b>Background</b>            | cytokine receptor like factor 3(CRLF3) Homo sapiens This gene encodes a cytokine receptor-like factor that may negatively regulate cell cycle progression at the G0/G1 phase. Studies of the related rat protein suggest that it may regulate neuronal morphology and synaptic vesicle biogenesis. This gene is one of several genes located in the neurofibromatosis type I tumor suppressor region on the q arm of chromosome 17, a region that is subject to microdeletions, duplications, chromosomal breaks and rearrangements. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 2 and 5. [provided by RefSeq, Aug 2012], |
| <b>Other</b>                 | CRLF3 CREME9 CYTOR4 P48, Cytokine receptor-like factor 3 (Cytokine receptor-like molecule 9) (CREME-9) (Cytokine receptor-related protein 4) (Type I cytokine receptor-like factor p48)  |

### Product Images:

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*