

IRP-1 (phospho Ser138) rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A16435	Rabbit	1 mg/ml	98399
Applications	IHC,IF,ELISA		
Reactivity	Human,Mouse,Rat		
Dilution	WB 1:500-2000 ,IHC: 1:100 - 1:300. IF: 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.		
Storage	-20°C/1 year		
Specificity	Phospho-IRP-1 (S138) Polyclonal Antibody detects endogenous levels of IRP-1 protein only when phosphorylated at S138.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	The antiserum was produced against synthesized peptide derived from human IREB1 around the phosphorylation site of Ser138. AA range:106-155		
Uniprot No	P21399		
Alternative names	ACO1; IREB1; Cytoplasmic aconitate hydratase; Aconitase; Citrate hydrolyase; Ferritin repressor protein; Iron regulatory protein 1; IRP1; Iron-responsive element-binding protein 1; IRE-BP 1		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	<p>aconitase 1(ACO1) Homo sapiens The protein encoded by this gene is a bifunctional, cytosolic protein that functions as an essential enzyme in the TCA cycle and interacts with mRNA to control the levels of iron inside cells. When cellular iron levels are high, this protein binds to a 4Fe-4S cluster and functions as an aconitase. Aconitases are iron-sulfur proteins that function to catalyze the conversion of citrate to isocitrate. When cellular iron levels are low, the protein binds to iron-responsive elements (IREs), which are stem-loop structures found in the 5' UTR of ferritin mRNA, and in the 3' UTR of transferrin receptor mRNA. When the protein binds to IRE, it results in repression of translation of ferritin mRNA, and inhibition of degradation of the otherwise rapidly degraded transferrin receptor mRNA. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct</p>		
Other	ACO1, Cytoplasmic aconitate hydratase		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
