

## LIMK-2 (phospho Ser283) rabbit pAb antibody

| Catalog No :                 | Source:   | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A17028                       | Rabbit  | 1 mg/ml         | 72232         |
| <b>Applications</b>          | WB,IHC,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat,Monkey  |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. ELISA: 1:5000. Not yet tested in other applications.  |                 |               |
| <b>Storage</b>               | -20°C/1 year  |                 |               |
| <b>Specificity</b>           | Phospho-LIMK-2 (S283) Polyclonal Antibody detects endogenous levels of LIMK-2 protein only when phosphorylated at S283.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human LIMK2 around the phosphorylation site of Ser283. AA range:249-298   |                 |               |
| <b>Uniprot No</b>            | P53671  |                 |               |
| <b>Alternative names</b>     | LIMK2; LIM domain kinase 2; LIMK-2  |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |                 |               |
| <b>Clonality</b>             | Polyclonal  |                 |               |
| <b>Isotype</b>               | IgG   |                 |               |
| <b>Conjugation</b>           |   |                 |               |
| <b>Background</b>            | <p>LIM domain kinase 2(LIMK2) Homo sapiens There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],</p> |                 |               |
| <b>Other</b>                 | LIMK2, LIM domain kinase 2  |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---